

Topological approach to multigranulation rough sets

Guoping Lin · Jiye Liang · Yuhua Qian

Received: date / Accepted: date

Abstract For further studying the theory of multigranulation rough sets (MGRS), we attempt to investigate a new theory on multigranulation rough sets from the topological view in this paper. We firstly explore multigranulation topological rough space and its topological properties by giving some new definitions and theorems. Then, topological granularity and topological entropy are proposed to characterize the uncertainty of the multigranulation topological rough space. Finally, based on the invariance of interior and closure operators of a target concept, a granulation selection algorithm is introduced to deal with the granularity selection issue in the multigranulation rough data analysis.

Keywords Rough sets · MGRS · Topology · Multigranulation topological rough space · Topological entropy

1 Introduction

Rough set theory (RS), originated by Pawlak [28], is a new mathematical tool to deal with incomplete, imprecise, and

uncertain knowledge. In the past decades, rough set theory has developed significantly due to its wide applications. Various generalized rough set models have been established and their corresponding properties or structures have been investigated intensively and extensively [1–4, 16, 39, 50, 52]. These extensional ways are mainly based on either relaxing an equivalence relation or generalizing partition to covering on the universe.

Topology is an important branch of mathematics aimed at studying the invariance of a given space under topological transformation (homeomorphism) [9], whose theories and applications have been studied in [6, 7, 14, 7, 17, 18]. In [29], Pawlak has pointed out that topology is closely related to rough set theory and convinced that topology structure of the rough set is one of key issues of rough set theory. The topology theory and rough set theory have been applied in many science and engineering areas such as in chemistry, biology, image processing, knowledge acquisition and pattern recognition. Therefore, how to combine rough set theory and topology theory becomes an interesting and natural research topic, in fact which has received considerable attention from the scholars in this community [21, 22, 12, 30, 31, 27, 36, 43, 49, 46, 51]. In particular, Skowron [41] and Wiweger [45] separately discussed this topic in 1988. Lin continued to discuss this topic and established a connection between fuzzy rough sets and topology [21]. Furthermore, by using the theories of the topology and the neighborhood systems, Polkowski [26] constructed and characterized topology spaces from rough sets based on information systems. In the literature [27], Polkowski pointed out: “topological aspects of rough set theory were recognized early in the framework of topology of partitions”. Lashin [15] generalized rough set theory in the frameworks of topology space and topologized information tables. Zhu [51] studied several covering-based rough sets on the topological view.

G. Lin

Key Laboratory of Computational Intelligence and Chinese Information Processing of Ministry of Education, School of Computer and Information Technology, Shanxi University, Taiyuan, 030006, Shanxi, China

Department of Mathematics and Information Science, Zhangzhou Normal University, Zhangzhou 363000, P. R. China

J. Liang

Key Laboratory of Computational Intelligence and Chinese Information Processing of Ministry of Education, School of Computer and Information Technology, Shanxi University, Taiyuan, 030006, Shanxi, China E-mail: ljy@sxu.edu.cn

Y. Qian

Key Laboratory of Computational Intelligence and Chinese Information Processing of Ministry of Education, School of Computer and Information Technology, Shanxi University, Taiyuan, 030006, Shanxi, China

Yang [44] investigated relationships among separation axioms and two topology spaces. Kortelainen [11] and Jarvinen [8] considered relationships among modified sets, topological spaces and rough sets based on a preorder. Qin [37] discussed the relationship between fuzzy rough set models and fuzzy topologies on a finite non-empty universe. Some other authors discussed relationships between generalized rough sets and the topology from different viewpoints (for example see [5, 10, 41]). Skowron et al. [41] generalized the classical approximation spaces to tolerance approximation spaces and discussed the problems of attribute reduction in these spaces. In addition, connections between fuzzy rough set theory and fuzzy topology were also investigated [16, 37, 40].

According to the different strategies, Qian and Liang [33] proposed multigranulation rough sets by employing multiple binary relations on the universe instead of a single one. A target concept was described through these granulations on the universe based on a user's different requirements or targets of problem solving. Since multigranulation rough set is an important research direction of the rough set theory, there have been many studies on this topic. For example, Liu et al. [24, 25] proposed covering fuzzy rough set based multigranulation rough sets. Xu et al. [42] investigated another generalized version, called variable precision multigranulation rough sets. Yang et al. [44] proposed a multigranulation rough set based on a fuzzy binary relation. Lin et al. [23] investigated a neighborhood-based multigranulation rough set model, which can be used to deal with the data sets with hybrid attributes. Liang et al. [20] proposed an efficient feature selection algorithm for large-scale data sets from the perspective of multigranulation which also demonstrates the usefulness of MGRS theory. What deserves to be mentioned is that She et al. [38] explored the topological structures and the properties of multigranulation rough sets. However, the multigranulation topological rough space and the corresponding topological properties are still not studied. The motivation of this paper is to investigate these problems and to describe the uncertainty of multigranulation topological space by proposing topological granularity measure and topological entropy.

With respect to the application of the multigranulation rough sets in the multi-source information system and the distributive information system where information is obtained from different sources [13], it is important to consider two interesting issues, i.e., the granular selection and the granularity selection in the view of granular computing [35]. They are two different important ways to reduce the redundant information in data analysis. The theory of granular selection is the same as that of covering reduct [53]. So we only concentrate on the issue of the granularity selection in this paper. Given two multigranulation spaces of a universe, two

multigranulation topological rough spaces will be generated. In the view of granular computing, the issue in this paper is that for a family of single granulation spaces which induced a multigranulation topological rough space, what would be the corresponding "smallest" subset of that family of granulations, which can produce the same multigranulation topological rough space? That is, they have the same interior and closure operators of a target concept.

Throughout the research, some new results and achievements proposed in the paper may enrich the theories of the multigranulation rough sets and the topology, which may form the theoretical basis for further applications of multigranulation rough set theory and topology.

The main objective of this paper is to study the multigranulation rough set theory via topology theory. The rest of this paper is organized as follows. Some basic concepts of topology and multigranulation rough sets are briefly reviewed in Section 2. In Section 3, the multigranulation topological rough space is constructed and some of its important properties are investigated. In Section 4, the topological granularity and the topological entropy are introduced to characterize the uncertainty of multigranulation topological rough space. The concept of granularity selection is proposed and a granularity selection algorithm based on the invariance of interior and closure operators of a target concept is given to select the necessary granulation in multigranulation rough data analysis. Finally, Section 5 concludes with some remarks.

2 Preliminaries

In this section, we introduce some fundamental key concepts of topology and rough set theory [29, 9]. Throughout this paper, we suppose that the universe Ω is a nonempty finite set.

2.1 Basic concepts of topology

We present a brief overview of topology space, a closure operator, an interior operator, and a topology based on a set. They are all important concepts in topological theory and they were used to study rough sets [22, 32, 47]. In this paper, these topological tools are also employed to investigate multigranulation rough sets.

Definition 2.1. (A topological space) [9] A topology space is a pair (Ω, τ) consisting of a set Ω and a family τ of subset of Ω satisfying the following conditions:

- (T1) $\emptyset \in \tau$ and $\Omega \in \tau$,
- (T2) τ is closed under arbitrary union,
- (T3) τ is closed under finite intersection.

The pair (Ω, τ) is called a topological space. The elements of Ω are called points of the space. The subsets of Ω belonging to τ are called open set in the space, and the complement of the subsets of Ω belonging to τ are called closed set in the space and the family of open subsets of Ω is also called a topology for Ω .

Definition 2.2. (Closure and interior operators). For an operator $cl: 2^\Omega \rightarrow 2^\Omega$, if it satisfies the following conditions, then we call it a closure operator on Ω . $\forall X, Y \subseteq \Omega$,

- (C1) $cl(\emptyset) = \emptyset$,
- (C2) $X \subseteq cl(X)$,
- (C3) $cl(cl(X)) = cl(X)$,
- (C4) $cl(X \cup Y) = cl(X) \cup cl(Y)$.

For an operator $int: 2^\Omega \rightarrow 2^\Omega$, if it satisfies the following rules, then we call it an interior operator on Ω . $\forall X, Y \subseteq \Omega$,

- (I1) $int(\Omega) = \Omega$,
- (I2) $int(X) \subseteq X$,
- (I3) $int(int(X)) = int(X)$,
- (I4) $int(X \cap Y) = int(X) \cap int(Y)$.

It is well known that an interior operator int on Ω can induce a topology τ such that in the topology space (Ω, τ) , $int(X)$ is just the interior of X for each $X \subseteq \Omega$. The similar statement is also true for a closure operator [31].

In a topology space (Ω, τ) , a family $\beta \subseteq \tau$ of sets is called a base for the topological τ if for each point x of the space, and each neighborhood X of x , there is a member V of β such that $x \in V \subseteq X$. We know that a subfamily β of a topology τ is a base for τ if and only if each member of τ is the union of members of β . Moreover, $\beta \subseteq 2^\Omega$ forms a base for some topology on Ω if and only if β satisfies the following conditions:

- (B1) $\Omega = \cup\{B|B \in \beta\}$,
- (B2) For every two members X and Y of β and each point $x \in X \cap Y$ there is $Z \in \beta$ such that $x \in Z \subseteq X \cap Y$.

2.2 Fundamentals of multigranulation rough sets

In [33], Qian analyzed some restrictions of Pawlak classical rough set in practice and proposed a new extension of rough set i.e., multigranulation rough sets, in which a target concept can be approximated by multiple equivalence relations according to a user's different requirements. In other words, a target concept can be approximated by multiple granularity spaces in the view of granular computing [34].

Assume that Ω is a finite nonempty universe of discourse. Let R be an equivalence relation on Ω , Ω/R is a corresponding partition of Ω , denoted by $\Omega/R = \{[x]_R|x \in \Omega\}$ in which $[x]_R = \{y|y \in \Omega, xRy\}$ is an equivalence class consisting x . Ω/R can generate a topology space, denoted as (Ω, τ_R) , and Ω/R is a topology base of τ_R , each subset of τ_R is both open and close [5].

Definition 2.3[33]. Let $S = (\Omega, AT, f)$ be an information system. Suppose that $X \subseteq \Omega$, R_1, R_2, \dots, R_q be q equivalence relations on Ω , the lower approximation $\underline{\sum_{i=1}^q R_i}(X)$ and the upper approximation $\overline{\sum_{i=1}^q R_i}(X)$ of X with respect to R_1, R_2, \dots, R_q are defined as follows, respectively,

$$\underline{\sum_{i=1}^q R_i}(X) = \{x \in \Omega \mid \forall ([x]_{R_i} \subseteq X), i \leq q\}; \quad (1)$$

$$\overline{\sum_{i=1}^q R_i}(X) = \{x \in \Omega \mid \wedge ([x]_{R_i} \cap X \neq \emptyset), i \leq q\}. \quad (2)$$

From the above expressions, the operator ' \forall ' is a disjunctive operator which here indicates that in multiple independent granular structures, one needs only at least one granular structure to satisfy with the inclusion condition between an equivalence class and a target concept. The expression (2) is the upper approximation of the optimistic multigranulation rough set that can be also defined by the complement of the lower approximation, which has been proved in [23]. the operator ' \wedge ' in expression (2) is a conjunctive operator whose meaning is that in multiple independent granular structures, one needs all granular structures to satisfy with non-empty for joint operator between an equivalence class and a target concept. And $\underline{\sum_{i=1}^q R_i}(X) \subseteq X \subseteq \overline{\sum_{i=1}^q R_i}(X)$. So we can label multigranulation rough set $X = (\underline{\sum_{i=1}^q R_i}(X), \overline{\sum_{i=1}^q R_i}(X))$, accordingly, we call $(\Omega, R_1, R_2, \dots, R_q)$ a multigranulation approximation space in the view of granular computing. From [33], we obtain the following interpretations:

- The lower approximation of a set X with respect to $\sum_{i=1}^q R_i$ is the set of all elements, which can certainly be classified as X using $\sum_{i=1}^q R_i$ (are certainly X in view of $\sum_{i=1}^q R_i$).
- The upper approximation of a set X with respect to $\sum_{i=1}^q R_i$ is the set of all elements, which can possibly be classified as X using $\sum_{i=1}^q R_i$ (are possibly X in view of $\sum_{i=1}^q R_i$).
- The boundary region of a set X with respect to $\sum_{i=1}^q R_i$ is the set of all elements, which can be classified neither as X nor as not- X using $\sum_{i=1}^q R_i$.

Let \emptyset be the empty set, $\sim X$ the complement of X in U , we have the following properties of multigranulation rough sets [33].

| | | |
|---|---------------------------------|---|
| (1ML) $\overline{\sum_{i=1}^q R_i(U)} = U$ | (Co-normality) | Proof. By Definition 3.1 and the definitions of lower and |
| (1MH) $\underline{\sum_{i=1}^q R_i(U)} = U$ | (Co-normality) | upper approximations of X in MGRS they can be easily |
| (2ML) $\overline{\sum_{i=1}^q R_i(\emptyset)} = \emptyset$ | (Normality) | proved. |
| (2MH) $\underline{\sum_{i=1}^q R_i(\emptyset)} = \emptyset$ | (Normality) | According to the above propositions of multigranulation |
| (3ML) $\overline{\sum_{i=1}^q R_i(X)} \subseteq X$ | (Contraction) | rough sets, we easily obtain the following results. |
| (3MH) $X \subseteq \underline{\sum_{i=1}^q R_i(X)}$ | (Extension) | Proposition 3.1. Let $(\Omega, \tau_1), (\Omega, \tau_2), \dots, (\Omega, \tau_q)$ be q topol- |
| (4ML) $\overline{\sum_{i=1}^q R_i(\bigcap_{j=1}^n X_j)} \subseteq \bigcap_{j=1}^n (\overline{\sum_{i=1}^q R_i(X_j)})$ | (Implication) | ogy spaces induced by equivalence relations R_1, R_2, \dots, R_q , |
| (4MH) $\underline{\sum_{i=1}^q R_i(\bigcup_{j=1}^n X_j)} \supseteq \bigcup_{j=1}^n (\underline{\sum_{i=1}^q R_i(X_j)})$ | (Implication) | respectively, and $X, Y \subseteq \Omega$. Then, with respect to the opera- |
| (5ML) $\overline{\sum_{i=1}^q R_i(\bigcup_{j=1}^n X_j)} \supseteq \bigcup_{j=1}^n (\overline{\sum_{i=1}^q R_i(X_j)})$ | (Implication) | tors <i>mint</i> , we have |
| (5MH) $\underline{\sum_{i=1}^q R_i(\bigcap_{j=1}^n X_j)} \subseteq \bigcap_{j=1}^n (\underline{\sum_{i=1}^q R_i(X_j)})$ | (Implication) | (1) $mint(\Omega) = \Omega$, |
| (6ML) $\overline{\sum_{i=1}^q R_i(\sum_{i=1}^q R_i(X))} = \overline{\sum_{i=1}^q R_i(X)}$ | (Idempotency) | (2) $mint(\emptyset) = \emptyset$, |
| (6MH) $\underline{\sum_{i=1}^q R_i(\sum_{i=1}^q R_i(X))} = \underline{\sum_{i=1}^q R_i(X)}$ | (Idempotency) | (3) $mint(X) \subseteq X$, |
| (7ML) $\overline{\sum_{i=1}^q R_i(\sim X)} = \sim \overline{\sum_{i=1}^q R_i(X)}$ | (Duality) | (4) $X \subseteq Y \Rightarrow mint(X) \subseteq mint(Y)$, |
| (7MH) $\underline{\sum_{i=1}^q R_i(\sim X)} = \sim \underline{\sum_{i=1}^q R_i(X)}$ | (Duality) | (5) $mint(mint(X)) = mint(X)$. |
| (8ML) $X \subseteq Y \Rightarrow \overline{\sum_{i=1}^q A_i(X)} \subseteq \overline{\sum_{i=1}^q R_i(Y)}$ | (Monotone) | Similarly, with respect to the operators <i>cl</i> , we have the |
| (8MH) $X \subseteq Y \Rightarrow \underline{\sum_{i=1}^q R_i(X)} \subseteq \underline{\sum_{i=1}^q R_i(Y)}$ | (Monotone) | following results: |
| (9ML) $\forall K \in U/R_i, i \in \{1, 2, \dots, q\}, \overline{\sum_{i=1}^q R_i(K)} = K$ | (Granularity) | (1) $mcl(\Omega) = \Omega$, |
| (9MH) $\forall K \in U/R_i, i \in \{1, 2, \dots, q\}, \underline{\sum_{i=1}^q R_i(K)} = K$ | (Granularity) | (2) $mcl(\emptyset) = \emptyset$, |
| (10ML) $\overline{\sum_{i=1}^q R_i(X)} = \bigcup_{i=1}^q (\overline{R_i(X)})$ | (Relation based Addition) | (3) $X \subseteq mcl(X)$, |
| (10MH) $\underline{\sum_{i=1}^q R_i(X)} = \bigcap_{i=1}^q (\underline{R_i(X)})$ | (Relation based Multiplication) | (4) $X \subseteq Y \Rightarrow mcl(X) \subseteq mcl(Y)$, |
| | | (5) $mcl(mcl(X)) = mcl(X)$. |

By the above discussions and similar to Definition 1 of [48], we can define the interior and closure operators of multigranulation rough set X from a topological point of view.

3 Topological approach to multigranulation rough sets

In this section, in order to better apply the multigranulation rough set theory into complex data analysis, we shall investigate an interesting and natural research topic of studying multigranulation rough set theory via topology theory.

Definition 3.1. Let $(\Omega, \tau_1), (\Omega, \tau_2), \dots, (\Omega, \tau_q)$ be q topological spaces induced by equivalence relations R_1, R_2, \dots, R_q , respectively, and $X \subseteq \Omega$. Then we define *mint*(X) and *mcl*(X) operators of X with respect to Γ , where $\Gamma = \{\tau_1, \tau_2, \dots, \tau_q\}$, respectively, as follows:

$$mint(X) = \bigcup \{A \in \tau_i \mid \forall (A \subseteq X), i \leq q\}; \quad (3)$$

$$mcl(X) = \bigcup \{A \in \tau_i \mid \forall (A \cap X \neq \emptyset), i \leq q\}. \quad (4)$$

The area of topological uncertainty or boundary (*mbn*) is defined as

$$mbn(X) = mcl(X) \setminus mint(X).$$

Theorem 3.1. Let $(\Omega, \tau_1), (\Omega, \tau_2), \dots, (\Omega, \tau_q)$ be q topological spaces induced by equivalence relations R_1, R_2, \dots, R_q , respectively, and $X \subseteq \Omega$. We have $mint(X) = \underline{\sum_{i=1}^q R_i(X)}$, $mcl(X) = \overline{\sum_{i=1}^q R_i(X)}$.

Theorem 3.2. Let $(\Omega, \tau_1), (\Omega, \tau_2), \dots, (\Omega, \tau_q)$ be q topological spaces induced by equivalence relations R_1, R_2, \dots, R_q , respectively, and $X, Y \subseteq \Omega$. Then, $mint(X \cap Y) = mint(X) \cap mint(Y)$ holds if and only if the multigranulation rough set model is equivalent to the Pawlak's single granulation rough set model.

Proof. It can be proved by employing the result of Proposition 10 in [20].

Theorem 3.3. Let $(\Omega, \tau_1), (\Omega, \tau_2), \dots, (\Omega, \tau_q)$ be q topological spaces induced by equivalence relations R_1, R_2, \dots, R_q , respectively, and $X, Y \subseteq \Omega$. Then, $mcl(X \cup Y) = mcl(X) \cup mcl(Y)$ holds if and only if the multigranulation rough set model is equivalent to the Pawlak's single granulation rough set model.

Proof. It can be proved by employing the result of Proposition 10 in [20].

From the above discussions, we can get the following results.

Theorem 3.4. Let $(\Omega, \tau_1), (\Omega, \tau_2), \dots, (\Omega, \tau_q)$ be q topological spaces induced by equivalence relations R_1, R_2, \dots, R_q , respectively, and $X, Y \subseteq \Omega$. Then, *mint*(X) and *mcl*(X) are interior and closure operators, respectively, if and only if the multigranulation rough set model is equivalent to the Pawlak's single granulation rough set model.

Definition 3.2. (Natural mapping). Let R be an equivalence relation, the family of $\{[x] \mid x \in \Omega\}$ is a quotient set on Ω , denoted by Ω/R . A mapping $f: \Omega \rightarrow \Omega/R$ satisfies $f(x) = [x], x \in \Omega$, we call f a natural mapping on Ω .

Definition 3.3. (Intersection operator). Let R_1 and R_2 be equivalence relations on a finite universe Ω , f_1 and f_2 natural mappings. Then we define an intersection mapping $F_{\cap} : \Omega \rightarrow 2^{\Omega}$ satisfies $F_{\cap}(x) = f_1(x) \cap f_2(x)$.

Further, if τ_1, τ_2 are two topologies induced by R_1 and R_2 , then we can define $\tau_1 \cap \tau_2 = \{F_{\cap}(x) \mid x \in \Omega\}$.

Theorem 3.5. $\tau_1 \cap \tau_2$ is a topology.

Proof. Suppose Ω be a finite universe, R_1 and R_2 two equivalence relations, and $\tau_1 = \{\emptyset, \Omega, [x_{i1}]_{R_1}, [x_{i2}]_{R_1}, \dots, [x_{ik}]_{R_1}\}$, $\tau_2 = \{\emptyset, \Omega, [x_{j1}]_{R_2}, [x_{j2}]_{R_2}, \dots, [x_{jl}]_{R_2}\}$ induced by R_1 and R_2 , $k, l \leq |\Omega|$, where $|\cdot|$ is cardinality of Ω .

(1) Based on the definition of $\tau_1 \cap \tau_2$, obviously $\emptyset \in \tau_1 \cap \tau_2$, $\Omega \in \tau_1 \cap \tau_2$.

(2) Assume that $X, Y \in \tau_1 \cap \tau_2$, then there exists two equivalence classes $[x]_{R_1} \in \tau_1, [x']_{R_2} \in \tau_2$ such that $X \subset [x]_{R_1}, Y \subset [x']_{R_2}$. Hence $X \cap Y \in [x]_{R_1} \cap [x']_{R_2} \in \tau_1 \cap \tau_2$.

(3) Let $\tau \in \tau_1 \cap \tau_2$, suppose that $\bigcup_{X \in \tau} X \notin \tau_1 \cap \tau_2$. Then there at least exists an element $x \in X \in \tau$, we have an equivalence class $[x]$ consisting x in $\tau_1 \cap \tau_2$ such that $[x] \notin \tau_1 \cap \tau_2$. Note that $[x] = ([x]_{R_1} \cap [x]_{R_2}) \in \tau_1 \cap \tau_2$ holds, a contraction!

Therefore, $\tau_1 \cap \tau_2$ is still a topology.

Similarly, we can prove that the intersection of the finite topologies is topology, i.e. $\prod_{i=1}^m \tau_i$ is a topology with respect to $\tau_1, \tau_2, \dots, \tau_q$, denoted by $\prod_{i=1}^m \tau_i = \Gamma_M$.

Definition 3.4. (Multigranulation topological rough space) Let $(\Omega, \tau_1), (\Omega, \tau_2), \dots, (\Omega, \tau_q)$ be q topology spaces induced by equivalence relations R_1, R_2, \dots, R_q , respectively. An intersection operation $F_{\cap}(x) : \Omega \rightarrow 2^{\Omega}$. Then $(\Omega, \prod_{i=1}^q \tau_i)$ is called a multigranulation topological rough space, denoted as $(\Omega, \prod_{i=1}^q \tau_i) = (\Omega, \Gamma_M)$, written by (Ω, Γ) for simplicity.

Definition 3.5. (The partial relation between two multigranulation topological rough space) Let τ_1, τ_2 be two topologies on Ω , if for any $X \in \tau_1$, there exists $Y \in \tau_2$ such that $X \subseteq Y$. Then we call τ_1 finer than τ_2 , denoted by $\tau_1 \leq^{\tau} \tau_2$. If τ_1 is strictly finer than τ_2 , denoted by $\tau_1 <^{\tau} \tau_2$. If and only if $X = Y$, then $\tau_1 = \tau_2$. Similarly, let Γ_1, Γ_2 be two multigranulation topological rough spaces on Ω , if for any $\tau_1 \in \Gamma_1$, there exists $\tau_2 \in \Gamma_2$ such that $\tau_1 \leq^{\tau} \tau_2$, then we call Γ_1 finer than Γ_2 , denoted by $\Gamma_1 \leq^{\Gamma} \Gamma_2$. If Γ_1 is strictly finer than Γ_2 , denoted by $\Gamma_1 <^{\Gamma} \Gamma_2$. If and only if $\tau_1 = \tau_1$, then $\Gamma_1 = \Gamma_2$.

Theorem 3.6. Let $\tau_1, \tau_2, \dots, \tau_q$ be q topologies on Ω induced by equivalence relations R_1, R_2, \dots, R_q , respectively. If $\tau_1 <^{\tau} \tau_2 <^{\tau} \dots <^{\tau} \tau_q$, then $\Gamma = \tau_1$.

Actually, from the above definition, we know (Ω, Γ) is finer than each topology on Ω .

Corollary 3.1. If $\tau_1 <^{\tau} \tau_2$, and β_1, β_2 are the topology base of τ_1, τ_2 , respectively. Then $\beta_1 <^{\tau} \beta_2$.

Corollary 3.2. If $\Gamma_1 <^{\Gamma} \Gamma_2$, and β_{1M}, β_{2M} are their family of the topology base of Γ_1, Γ_2 , respectively. Then $\beta_{1M} <^{\Gamma} \beta_{2M}$.

Theorem 3.7. (Topology base of Γ_M) Let β_i be the topology base of τ_i , then $\prod_{i=1}^q \beta_i$ is a topology base of multigranulation topological rough space Γ .

Proof. (1) For any $x \in \Omega$, there exists $[x]_{R_i} \in \beta_i$ such that $x \in \cap [x]_{R_i}$. Note that $\cap [x]_{R_i} \in \prod_{i=1}^q \beta_i$. Hence, let $B = \cap [x]_{R_i} \in \prod_{i=1}^q \beta_i$, we have $x \in B$.

(2) For any $B_1, B_2 \in \prod_{i=1}^q \beta_i$, suppose $x \in B_1 \cap B_2$, but B_1 is a set which is intersection of $\cap_{i=1}^q [x]_{R_i}$, B_2 is a set which is intersection of $\cap_{i=1}^q [y]_{R_i}$, then $x \neq y$, otherwise $[x]_{R_i} = [y]_{R_i}$. Hence $B_1 \cap B_2 = \emptyset$. There exists $B_3 = \emptyset$, obviously, $\emptyset \subset \emptyset$ holds.

Therefore $\prod_{i=1}^q \beta_i$ is a topology base of multigranulation topological rough space Γ .

Corollary 3.3. Let $\tau_1, \tau_2, \dots, \tau_q$ be q topologies on Ω induced by equivalence relations R_1, R_2, \dots, R_q and β_i the topology base of τ_i for $i \in \{1, 2, \dots, q\}$. Then $\prod_{i=1}^q \beta_i$ is a partition of Ω .

Definition 3.5. Let (Ω, Γ) be a multigranulation topological rough space, β_M is a topology base of Γ , and $X \subseteq \Omega$. Then the interior operator is defined as

$$INT(X) = \{x \in X \mid \forall x \in A \in (\Gamma \setminus N), \text{ then } A \subseteq X\}, \quad (5)$$

where $N = \bigcup \{Y \mid Y \subseteq X \subseteq \Omega\}$.

Definition 3.6. Let (Ω, Γ) be a multigranulation topological rough space, β_M a topology base of Γ , and $X \subseteq \Omega$. Then the closure operator is

$$CL(X) = \bigcup \{A \in \Gamma \mid A \cap X \neq \emptyset\}. \quad (6)$$

Example 3.1. Let $\Omega = \{x_1, x_2, x_3, x_4, x_5\}$ and $X = \{x_1, x_2, x_3, x_4\} \subseteq \Omega$. R_1 and R_2 are two equivalence relations on Ω . $\Omega/R_1 = \{\{x_1, x_2, x_3\}, \{x_4, x_5\}\}$ corresponding to $f_1(x) = [x]_{R_1}$, $\Omega/R_2 = \{\{x_1\}, \{x_3\}, \{x_2, x_4, x_5\}\}$ corresponding to $f_2(x) = [x]_{R_2}$. Then we have that $\sum_{i=1}^2 R_i(X) = \{x_1, x_2, x_3\}$, $\sum_{i=1}^2 R_i(X) = \{x_1, x_2, x_3, x_4, x_5\}$ and get two topologies $\tau_1 = \{\emptyset, \Omega, \{x_1, x_2, x_3\}, \{x_4, x_5\}\}$, $\tau_2 = \{\emptyset, \Omega, \{x_1\}, \{x_3\}, \{x_2, x_4, x_5\}\}$. $\beta_1 = \{\{x_1, x_2, x_3\}, \{x_4, x_5\}\}$ and $\beta_2 = \{\{x_1\}, \{x_3\}, \{x_2, x_4, x_5\}\}$ are topology bases of τ_1 and τ_2 , respectively. By Definition 3.3, we have that $\Gamma = \tau_1 \cap \tau_2 = \{\emptyset, \Omega, \{x_1, x_2, x_3\}, \{x_4, x_5\}, \{x_1\}, \{x_3\}, \{x_2, x_4, x_5\}, \{x_1, x_3\}, \{x_1, x_2\}, \{x_2, x_3\}, \{x_2, x_3, x_4, x_5\}, \{x_1, x_2, x_4, x_5\}\}$. Then the topology base of $\beta_M = \{\{x_1\}, \{x_3\}, \{x_2\}, \{x_4, x_5\}\}$. Hence, $mint(X) = \{x_1, x_2, x_3\}$, $mcl = \{x_1, x_2, x_3, x_4, x_5\}$.

As a result of this example, we have the following propositions.

Proposition 3.1. Let (Ω, Γ) be a multigranulation topological rough space and $X \subseteq \Omega$. $\sum_{i=1}^q R_i(X)$ and $\overline{\sum_{i=1}^q R_i(X)}$ are lower and upper approximations of X . Then we have

$$(1) \underline{\sum_{i=1}^q R_i X} = INT(X),$$

$$(2) \overline{\sum_{i=1}^q R_i X} = CL(X).$$

Proof. (1) For any $x \in \sum_{i=1}^q R_i X$, there exists $[x]_R \subseteq X$. From Definition 3.4, $[x]_R \in \overline{\Gamma_M} \setminus N$, where $N = \{Y | Y \subseteq X\}$. Hence $x \in INT(X)$, i. e., $\sum_{i=1}^q R_i X \subseteq INT(X)$. On the other hand, for any $x \in INT(X)$, there exists $A \in \Gamma \setminus N$ such that $x \in A$ and $A \subseteq X$. Obviously, A is an equivalence class induced by R and $A = [x]_R$, i. e., $[x]_R \subseteq X$. So $x \in \sum_{i=1}^q R_i X$. Hence $x \in INT(X) \subseteq \sum_{i=1}^q R_i X$. Therefore, $\sum_{i=1}^q R_i X = INT(X)$ holds.

(2) According to the definition of $\sum_{i=1}^q R_i X$, for any R_i , $[x]_{R_i} \cap X \neq \emptyset$ holds. To Γ , there exists some element V (a subset of Ω) in Γ such that $V \subseteq [x]_{R_i}$. So there exists two cases:

(i) If $V \cap X = \emptyset$, then $V \not\subseteq CL(X)$. Hence $\overline{\sum_{i=1}^q R_i X} = CL(X)$ holds.

(ii) If $V \cap X \neq \emptyset$, then $V \subseteq CL(X)$. Note that $V \subseteq [x]_{R_i}$, hence $\overline{\sum_{i=1}^q R_i X} = CL(X)$ also holds.

From the above Proposition, we find that a target concept's interior and closure operators in multigranulation topological rough space are also equal to its lower and upper approximations in the multigranulation rough sets, respectively.

Proposition 3.2. Let (Ω, Γ) be a multigranulation topological rough space and $X \subseteq Y \subseteq \Omega$. Then $INT(X) \subseteq INT(Y)$, $CL(X) \subseteq CL(Y)$.

Proof. They can be proved similar to Theorem 3.2 in [34].

4 Measure of multigranulation topological rough space

4.1 Measure of multigranulation topological rough space

In this section, we introduce the uncertainty of multigranulation topological rough space.

Definition 4.1. (Granularity of a set) Let Ω be a finite nonempty universe. A function $m : 2^\Omega \rightarrow \mathfrak{R}$ is called a measure of the granularity of a set if it satisfies the following conditions: for all $A, B \in 2^\Omega$,

$$(M1) \quad m(A) \geq 0,$$

$$(M2) \quad A \subset B \Rightarrow m(A) < m(B),$$

$$(M3) \quad A \sim_s B \Rightarrow m(A) = m(B).$$

Where $A \sim_s B \Leftrightarrow (\sim(A <_s B), \sim(B <_s A))$, " $<_s$ " is the weak order that is an extension of " \subset " (see [47]).

Theorem 4.1. Let $\Omega/R = \{A_1, A_2, \dots, A_k\}$ be a partition of Ω , then we call

$$m(A) = \frac{|\Omega|}{k} \left(1 - \frac{1}{k \cdot |A|}\right)$$

a measure of granularity of a set A , k is the number of blocks in Ω/R , denoted by $|\Omega/R| = k$.

Proof. It is sufficient to show that m meets all the conditions in Definition 4.1.

$$(1) \text{ Obviously, } m(A) = \frac{|\Omega|}{k} \left(1 - \frac{1}{k \cdot |A|}\right) \geq 0.$$

$$(2) \text{ If } A \subset B, \text{ then } |A| < |B|, \text{ then } m(A) - m(B) = \frac{|\Omega|}{k} \left(1 - \frac{1}{k \cdot |A|}\right) - \frac{|\Omega|}{k} \left(1 - \frac{1}{k \cdot |B|}\right) = \frac{|\Omega|}{k} \left(1 - \frac{1}{k \cdot |A|} - 1 + \frac{1}{k \cdot |B|}\right) = \frac{|\Omega|}{k} \left(\frac{1}{k \cdot |B|} - \frac{1}{k \cdot |A|}\right) < 0, \text{ i.e., } m(A) < m(B).$$

$$(3) \text{ If } A \sim_s B, \text{ then } |A| \leq |B| \text{ and } |A| \geq |B|. \text{ Hence } m(A) = m(B).$$

Proposition 4.1. (Maximum) Let Ω/R be a partition of Ω induced by an equivalence relation on Ω and $X \in \Omega/R$. The maximum granularity measure of X with respect to R is one. This value is achieved if and only if $k = 1$, $\max(m(A)) = |\Omega| \left(1 - \frac{1}{|\Omega|}\right)$.

Proposition 4.2. (Minimum) Let Ω/R be a partition of Ω induced by an equivalence relation on Ω and $X \in \Omega/R$. The minimum granularity measure of X with respect to R is one. This value is achieved if and only if $k = |\Omega|$, $\min(m(A)) = \left(1 - \frac{1}{|\Omega|}\right)$.

$$\text{Hence, } \left(1 - \frac{1}{|\Omega|}\right) \leq m(A) \leq |\Omega| \left(1 - \frac{1}{|\Omega|}\right).$$

Example 4.1 (Continued from Example 3.1) To β_1 , $m(B_{11}) = \frac{5}{2} \left(1 - \frac{1}{3 \cdot 2}\right) = \frac{25}{12}$, $m(B_{12}) = \frac{5}{2} \left(1 - \frac{1}{2 \cdot 2}\right) = \frac{15}{8}$, so $m(\beta_1) = \left(\frac{25}{12}, \frac{15}{8}\right)^T$. To β_2 , $m(B_{21}) = \frac{2}{9}$, $m(B_{22}) = \frac{2}{9}$, $m(B_{23}) = \frac{8}{27}$, so $m(\beta_2) = \left(\frac{2}{9}, \frac{2}{9}, \frac{8}{27}\right)^T$. To β_M , $m(B_1) = \frac{4}{5}$, $m(B_2) = \frac{4}{5}$, $m(B_3) = \frac{9}{10}$, $m(B_4) = \frac{9}{10}$, $m(B_4) = \frac{9}{10}$, so $m(\beta_M) = \left(\frac{4}{5}, \frac{4}{5}, \frac{9}{10}, \frac{9}{10}, \frac{9}{10}\right)^T$.

Definition 4.2. Let $T = \{\Gamma\}$ be a family of multigranulation topological rough spaces on Ω . A function $G : \Gamma \rightarrow \mathfrak{R}$ is called a measure of granularity of a partition if it satisfies the following conditions for all $\Gamma_1, \Gamma_2 \in T$,

$$(G1) \quad G(\Gamma) \geq 0 \quad (\text{Nonnegativity})$$

$$(G2) \quad \Gamma_1 \subset \Gamma_2 \Rightarrow G(\Gamma_1) < G(\Gamma_2) \quad (\text{Monotonicity})$$

$$(G3) \quad \Gamma_1 = \Gamma_2 \Rightarrow G(\Gamma_1) = G(\Gamma_2) \quad (\text{Size invariance})$$

Considering a family set $\beta_M = \{B_1, B_2, \dots, B_q\}$ of a finite nonempty universe Ω . One may associate it with a probability discussion [19], $P_{\beta_M} = \left(\frac{|B_1|}{|\Omega|}, \frac{|B_2|}{|\Omega|}, \dots, \frac{|B_q|}{|\Omega|}\right)$. Then we can define granularity of a multigranulation topological rough space as follows.

Theorem 4.2. (Topological granularity) Let (Ω, Γ) be a multigranulation topological rough space, $m : 2^\Omega \rightarrow \mathfrak{R}$ a measure of the granularity of subsets of Ω , and $\beta_M = \{B_1, B_2, \dots, B_q\}$ a topology base of Γ_M . Then a measure

$$G_M(\Gamma) = \sum_{i=1}^q m(B_i) \cdot p(B_i).$$

is topological granularity of Γ , where $p(B_i) = \frac{|B_i|}{|\Omega|}$.

Proof. It is sufficient to show that G_M satisfies all the conditions in Definition 4.2.

$$(1) \text{ Obviously, } G_M(\Gamma) \geq 0 \text{ holds.}$$

(2) Suppose $\Gamma_1 <^\Gamma \Gamma_2$, by Corollary 3.3, $\beta_{1M} \subset \beta_{2M}$ holds. This means that every equivalence class of β_{2M} is a

union of one or more blocks of β_{1M} and at least one equivalence class of β'_M is the union of at least two blocks from β_{1M} . By the fact Ω is a finite universe, there exists a finite sequence of partitions $\beta_{1M} = \beta_{M1} \subset \beta_{M2} \subset \dots \subset \beta_{Ml} = \beta_{2M}$ such that exactly one block of β_{j+1} is the union of two equivalence classes from β_j for $j = 1, 2, \dots, n-1$ and $n \geq 2$. We want to show that $G(\beta_j) < G(\beta_{j+1})$. Without loss of generality, suppose a equivalence class of β_{j+1} is obtained by the union of two equivalence classes B_{j1} and B_{j2} of β_j , that is, $\beta_j = \{B_{j1}, B_{j2}, \dots, B_{jk}\}, k \geq 2$ and $\beta_{j+1} = \{B_{j1} \cup B_{j2}, \dots, \cup B_{jk}\}$. According to the definition of $G_M(\Gamma)$ and monotonicity of m , we have:

$$\begin{aligned} G_M(\beta_j) &= \sum_{i=1}^k m(B_{ji}) \cdot p(B_{ji}) \\ &= m(B_{j1}) \cdot p(B_{j1}) + m(B_{j2}) \cdot p(B_{j2}) + \sum_{i=3}^k m(B_{ji}) \cdot p(B_{ji}) \\ &< m(B_{j1} \cup B_{j2}) \cdot p(B_{j1}) + m(B_{j2} \cup B_{j1}) \cdot p(B_{j2}) + \sum_{i=3}^k m(B_{ji}) \cdot p(B_{ji}) \\ &= m(B_{j1} \cup B_{j2}) \cdot (p(B_{j1}) + p(B_{j2})) + \sum_{i=3}^k m(B_{ji}) \cdot p(B_{ji}) \\ &= m(B_{j1} \cup B_{j2}) \cdot p(B_{j1} \cup B_{j2}) + \sum_{i=3}^k m(B_{ji}) \cdot p(B_{ji}) \\ &= G_M(\beta_{j+1}). \end{aligned}$$

It immediately follows that $G(\Gamma_1) < G(\Gamma_2)$ holds.

(3) Suppose $\Gamma_1 = \Gamma_2$, based on Corollary 3.3, $\beta_{1M} \subset \beta_{2M}$ holds. And by Definition 3.4, $G(\Gamma_1) = G(\Gamma_2)$ holds.

Proposition 4.3. (Maximum). Let (Ω, Γ) be a multigranulation topological rough space. The maximum topology granularity measure of τ with respect to Ω is one. This value is achieved if and only if $m = 1$, $\max(G_M(\Gamma)) = |\Omega| - 1$.

Proposition 4.4. (Minimum). Let (Ω, Γ) be a multigranulation topological rough space. The maximum topology granularity of τ with respect to Ω is one. This value is achieved if and only if $m = |\Omega|$, $\min(G_M(\Gamma)) = 1 - \frac{1}{|\Omega|}$.

$$\text{Hence, } 1 - \frac{1}{|\Omega|} \leq G_M(\Gamma) \leq |\Omega| - 1.$$

Theorem 4.3. Let Γ_1, Γ_2 be two multigranulation topological rough space. If $\Gamma_1 <^\tau \Gamma_2$, then $G_M(\Gamma_1) < G_M(\Gamma_2)$.

Theorem 4.4. Let $\tau_1, \tau_2, \dots, \tau_q$ be q topologies induced by equivalence relations R_1, R_2, \dots, R_q , respectively. If $\tau_1 <^\tau \tau_2 <^\tau \dots <^\tau \tau_q$, then $G_M(\Gamma) = G_M(\tau_q)$.

Definition 4.3. (Topological entropy) Let (Ω, Γ) be a multigranulation topological rough space, $\beta_M = \{B_1, B_2, \dots, B_q\}$ is a topology base of Γ . Then the topological entropy of Γ is defined as:

$$E_M(\Gamma) = 1 - \frac{1}{|\Omega|} \sum_{i=1}^q m(B_i) \cdot p(B_i),$$

$$\text{where } p(B_i) = \frac{|B_i|}{|\Omega|}.$$

Proposition 4.5. (Maximum). Let (Ω, Γ) be a multigranulation topological rough space. The maximum topology entropy of τ with respect to Ω is one. This value is achieved if and only if $q = |\Omega|$, $\max(E_M(\Gamma)) = 1 - \frac{1}{|\Omega|} + \frac{1}{|\Omega|^2} = 1$.

Proposition 4.6. (Minimum) Let (Ω, Γ) be a multigranulation topological rough space. The maximum topology entropy measure of τ with respect to Ω is one. This value is achieved only if $q = 1$, $\min(E_M(\Gamma)) = \frac{1}{|\Omega|}$.

$$\text{Hence, } \frac{1}{|\Omega|} \leq E_M(\Gamma) \leq 1 - \frac{1}{|\Omega|} + \frac{1}{|\Omega|^2}.$$

Example 4.2. (Continued from Example 3.1) From Example 4.1, $P_\Gamma = \{\frac{1}{5}, \frac{1}{5}, \frac{1}{5}, \frac{2}{5}\}$, then we have $G_M(\Gamma) = \frac{22}{25}$ and $E_M(\Gamma) = \frac{103}{125}$.

4.2 Application of multigranulation topological rough space

Multigranulation rough set model is one of important extensions of Pawlak rough set model. One of the advantage of the former may be suitable to deal with the complex problem, such as multi-source information system where information comes from different sources. In such multi-source environment, the granular selection and the granularity selection are two key issues in process of the multigranulation rough data analysis. Granular selection theory is the same as covering reduct theory. Accordingly, we only investigate the granularity selection theory in this section.

In what follows, we give a real-life example to illustrate the application of the multigranulation rough set theory via topology theory. For example, when a doctor will incline to diagnose a disease of a patient more accurately, he always integrates multiple values of the patient's physical examination indicators from different hospitals where supply different examination indicators. These integrated information coming from different sources is called the multi-source information.

Example 4.3. Let $U = \{x_1, x_2, x_3, x_4, x_5\}$ be a universe of six objects which are here regarded as patients. Suppose there are three hospitals($H_i, i = \{1, 2, 3, 4\}$) providing us information regarding the attributes $\{a, b, c\}, \{a, b, d\}, \{a, b, e\}, \{f, g, h\}$, respectively. These attributes represent the patient's physical examination indicators. Table 1 depicts the information provided by the three hospitals. In Table 1, a denotes $HBsAg$, b denotes $RHBs$; c denotes $HBeAg$, d denotes $RHBe$, e denotes $RHBc$, f denotes CHO , g denotes TG , and h denotes TP which are the eight examination indicators used to determine whether the patient has the disease of *Hepatitis B* or not. "+" and "-" represent two attribute values that the former indicates *positive* and the latter indicates *negative*. "H" "L" and "M" represent three attribute values that "H" indicates *High*, "L" indicates *Low*, and "M" indicates *Middle*. The data in Table 1 come from the URL:

<http://www.forwardhealth.wi.gov/.../PEHIUserGuide.pdf.spage>.

According to each subsystem where information was provided by each hospital can generate a granulation in the view

of granular computing. Accordingly, this multi-source information system may generate three granulations. However, these granulations are not equally significant or even some of those granulations are redundant that lead to too much cost for one patient. Seen by this way, it is necessary to delete some redundant granulations in the process of multi-granulation rough data analysis.

Definition 4.2. (Significance of granularity) We say that τ_k is significant in Γ , if $E_\Gamma(\Omega, \sqcap_{i=1}^q \tau_i) \neq E_\Gamma(\Omega, \sqcap_{i=1, i \neq k}^q \tau_i)$.

Whereas, τ_k is not significant in Γ , if $E_\Gamma(\Omega, \sqcap_{i=1}^q \tau_i) = E_\Gamma(\Omega, \sqcap_{i=1, i \neq k}^q \tau_i)$.

To further study significance of τ_k , we introduce a quantitative measure for the significance as follows.

The significance measure of τ_k in Γ is defined as

$$S_\Gamma(\tau_k) = E_\Gamma(\Omega, \sqcap_{i=1}^q \tau_i) \setminus E_\Gamma(\Omega, \sqcap_{i=1, i \neq k}^q \tau_i).$$

Example 4.4. (Continued from Example 3.1) From Example 4.1, we have $S_\Gamma(\tau_1) = \frac{-414}{3375} < 0$, $S_\Gamma(\tau_2) = \frac{6}{15}$.

Definition 4.3. (Granularity selection) Let $\Gamma = \{\tau_1, \tau_2, \dots, \tau_q\}$ be a family of topological spaces on Ω , if there exists a subset $\Gamma_i = \{\tau_{i1}, \tau_{i2}, \dots, \tau_{ik}\} \subseteq \Gamma$, such that $E_\Gamma(\Omega, \sqcap_{i=1}^q \tau_i) = E_\Gamma(\Omega, \sqcap_{i=k}^q \tau_{ik})$, but $E_\Gamma(\Omega, \tau_{i1} \sqcap \tau_{i2} \sqcap \dots \sqcap \tau_{ik} \sqcap \tau_{i(k+1)}) \neq E_\Gamma(\Omega, \tau_{i1} \sqcap \tau_{i2} \sqcap \dots \sqcap \tau_{iq})$, then we call Γ_i a granularity reduct of Γ .

Algorithm 4.1 (Granularity selection algorithm)

Input: A MSIDT $\mu = (U, (\{R_i\})_{i \in N})$, where MSIDT denotes a multi-source information decision table

Output: One reduct *reduct*.

Steps are shown as follows:

I1: *reduct* $\leftarrow \emptyset$; // *reduct* is the set to conserve the selected granularities

I2: For($i = j$; $j \leq |R_i|$; $j++$) Do

 Compute $S_\Gamma(\tau_k)$, $k \leq |R_i|$;

 Put τ_k into *reduct*, where $S_\Gamma(\tau_k) > 0$; // These granularities form the core of the given multi-source information decision table

 While $E_\Gamma(\Omega, \text{reduct}) = E_\Gamma(\Omega, \sqcap_{i=1}^q \tau_i)$,

 Break;

 Endfor

I3: Return *reduct*.

Step *I2* is one of the key steps in this algorithm. The time complexity of computing is $O(m|R_i||U|^2)$, where m is the number of the attribute of each granularity R_i . $|R_i|$ is the number of granularities on U . $|U|$ is the number of the samples on U .

According the above algorithm, we can get one granularity selection $\{H_1, H_2, H_3\}$ which can determine whether one suffers from the disease of *Hepatitis B* from Table 1, which

Table 2 A complete target information system about emporium investment project

| Project | Locus | Investment | Population density | Decision |
|---------|--------|------------|--------------------|----------|
| x_1 | common | high | big | yes |
| x_2 | bad | high | big | no |
| x_3 | bad | low | small | no |
| x_4 | bad | low | small | yes |
| x_5 | bad | low | small | no |

result is consistent to that determined by doctors in the real patient cases.

Example 4.5 Here, we employ another simple example to illustrate the effectiveness of the granularity selection algorithm. Table 2 depicts a complete target information system containing some information about an emporium investment project. *Locus*, *Investment* and *Population density* are the conditional attributes of the system, whereas *Decision* is the decision attribute. The attribute domains are as follows:

$V_{Locus} = \{good, common, bad\}$, $V_{Investment} = \{high, low\}$, $V_{Populationdensity} = \{big, small, medium\}$, $V_{Decision} = \{yes, no\}$.

Each attribute in Table 2 here is regarded as a granulation as well as a source, hence, we call it a special multi-source information system.

Based on Table 2, $\Omega = \{x_1, x_2, x_3, x_4, x_5\}$, suppose a target concept $X = \{x_2, x_4, x_5\} \subseteq \Omega$, by employing the multi-granulation rough set theory and multigranularity topological rough space theory, we can get all topological spaces as follows:

$\tau_1 = \{\emptyset, \Omega, \{x_1\}, \{x_2, x_3, x_4, x_5\}\}$,

$\tau_2 = \{\emptyset, \Omega, \{x_1, x_2\}, \{x_3, x_4, x_5\}\}$,

$\tau_3 = \{\emptyset, \Omega, \{x_1, x_2\}, \{x_3, x_4, x_5\}\}$,

$\tau_4 = \{\emptyset, \Omega, \{x_1\}, \{x_2\}, \{x_3, x_4, x_5\}\}$,

$\tau_5 = \{\emptyset, \Omega, \{x_1, x_2\}, \{x_3, x_4, x_5\}\}$,

$\tau_6 = \{\emptyset, \Omega, \{x_1\}, \{x_2\}, \{x_3, x_4, x_5\}\}$.

By Definition 3.4, then we can get a multigranulation topological rough space (Ω, Γ) , where $\Gamma = \{\emptyset, \Omega, \{x_1\}, \{x_2\}, \{x_3, x_4, x_5\}, \{x_1, x_2\}, \{x_1, x_3, x_4, x_5\}, \{x_2, x_3, x_4, x_5\}\}$. By the granularity selection algorithm, we know $\tau_1, \tau_2, \tau_3, \tau_4, \tau_5$ are redundant granularity spaces, and $\{\tau_6\}$ is a granularity selection of Γ which preserves the invariance of topological entropy of X in the multigranulation topological rough space.

5 Conclusions and discussions

In this paper, we have presented an investigation of the topology method for multigranulation rough sets and addressed a series of concepts of the multigranulation topological rough space and its topological properties. Moreover, we have introduced topological granularity and topological entropy to show the uncertainty of multigranulation topological rough

Table 1 A multi-source information system

| | H_1 | | | H_2 | | | H_3 | | | H_4 | | |
|---|-------|-----|-----|-------|-----|-----|-------|-----|-----|-------|-----|-----|
| | a | b | c | a | b | d | a | b | e | f | g | h |
| 1 | + | - | + | + | + | - | + | - | + | H | H | L |
| 2 | + | + | + | - | - | + | - | + | + | L | M | L |
| 3 | + | - | - | - | - | + | - | - | + | H | L | H |
| 4 | - | + | - | + | - | + | - | - | + | L | L | H |
| 5 | - | - | - | - | - | + | - | - | + | H | M | M |
| 6 | - | + | - | - | - | - | - | - | - | H | H | M |
| 7 | - | - | - | - | - | - | - | - | + | M | M | L |

space from the topological view. In particular, a granularity selection algorithm based on the invariance of topological entropy was preliminarily proposed to reduce redundant granulations in the multigranulation rough data analysis.

Acknowledgements The authors would like to thank the anonymous reviewers and the editor for their constructive and valuable comments. This work is supported by grants from National Natural Science Foundation of China under Grant (Nos. 60903110, 71140004, 11061004), National Key Basic Research and Development Program of China (973) (No. 2007CB311002), the Construction Project of the Science and Technology Basic Condition Platform of Shanxi(No.2012091002-0101), and Education Committee of Fujian Province under Grant (Nos. JK2011031).

References

- Abo-Tabl EA (2012) Rough sets and topological spaces based on similarity. *J Mach Learn Cybern* DOI 10.1007/s13042-012-0107-7.
- Bonikowski Z, Brynirski E, Wybraniec. U (1998) Extensions and intentions in the rough set theory. *Inf Sci* 107:149-167.
- Chen JK, Li JJ, Lin YJ (2012) On the structure of definable sets in covering approximation spaces. *J Mach Learn Cybern*. DOI 10.1007/s13042-012-0086-8.
- Chen DG, Wang CZ, Hu QH (2007) A new approach to attribute reduction of consistent and inconsistent covering decision systems with covering rough sets. *Inf Sci* 177: 3500-3518.
- Chen DG, Zhang WX (2001) Rough sets and topology space. *J Xi'an Jiaotong University* 35 (12):1313-1315 (in Chinese).
- Ciobanu G, Rusa D(2006) A topological approach of the web classification. *Lect Notes Comput Sci* 4281:80-92.
- Fang J (2007) L-fuzzy Alexandrov topologies and specialization orders. *Fuzzy Sets Syst* 158:2359-2374.
- Jarvinen J, Kortelainen J (2007) A unifying study between model-like operators, topologies, and fuzzy sets. *Fuzzy Sets Syst* 158:1217-1225.
- Kelley J (1995) General topology. Van Nostrand Company.
- Kondo M (2006) On the structure of generalized rough sets. *Inf Sci* 176:589-600.
- Kortelainen J (1994) On relationship between modified sets, topology spaces and rough sets. *Fuzzy Sets Syst* 61:91-95.
- Kang XP, Li DY (2012) Dependency space, closure system and rough set theory. *International J Mach Learn Cybern*. DOI 10.1007/s13042-012-0106-8.
- Khan Md.A, Banerjee M (2008) Formal reasoning with rough sets in multiple-source approximation systems. *Int J Approx Reason* 49:466-477.
- Lai H, Zhang D (2006) Fuzzy preorder and fuzzy topology. *Fuzzy Sets Syst* 157:1865-1885.
- Lashin EF, Kozae AM, Abo-Khadra AA, Medhat T (2005) Rough set theory for topology spaces. *Int J Approx Reason* 40:35-43.
- Li JH, Mei CL, Lv YJ (2012) Knowledge reduction in real decision formal contexts, *Inf Sci* 189:191-207
- Pei D, Xu Z (2007) Transformation of rough set models. *Knowl Based Syst* 20(8):745-751.
- Rasiowa H (1974) *An Algebraic Approach to Non-classical Logic*. North-Holland, Amsterdam.
- Lee TT (1987) An information-theoretic analysis of relational database, part I: data dependencies and information metric. *IEEE Trans Soft Engin* 13:1049-1061.
- Liang JY, Wang F, Dang CY, Qian YH (2012) An efficient rough feature selection algorithm with a multi-granulation view, *Int J Approx Reason* 53(7):1080-1093.
- Lin TY (1992) Topological and fuzzy rough sets in Decision Support by Experience-Application of the Rough Sets Theory, R. Slowinski, Ed. Norwell, MA: Kluwer, pp:287-304.
- Lin TY, Liu Q, Yao YY (1994) Logic systems for approximate reasoning: Via rough sets and topology in Methodologies for Intelligent Systems. Berlin, Germany: Springer-Verlag.
- Lin GP, Qian YH, Li JJ (2012) NMGRS: Neighborhood-based multigranulation rough sets, *Int J Approx Reason* 53(7) 1080-1093.
- Liu CH, Wang MZ (2011) Covering fuzzy rough set based on multi-granulations, *Int Conf Uncertainty Reason Knowl Engineering* (146-149).
- Liu CH, Miao DQ (2011) Covering rough set model based on multigranulations, *RSFDGRc2011* 87-90.
- Polkowski L (2001) On fractals defined in information systems via rough set theory. In: *Proceedings of the RSTGC-2001*, Bull. Int. Rough Set Society 5:163-166.
- Polkowski L, Semeniuk-Polkowska M (2004) Some remarks on sets of communicating sequential processes in topological rough set framework. *Fundam. Inform.* 60:291-305.
- Pawlak Z (1982) Rough sets. *Int J Comput Inf Sci* 5:341-356.
- Pawlak Z (1991) Rough sets: theoretical aspects of reasoning about data. Kluwer, Dordrecht.
- Pei D, Xu Z (2004) Rough set models on two universes. *Int J Gen Syst* 332 (5):569-581.
- Pei Z, Pei. DW, Zheng L (2011) Topology vs generalized rough sets. *Int J Approx Reason* 52:231-239.
- Pomykala JA (1987) Approximation operations in approximation space. *Bull Polish Academic Sci* 35 (9-10):653-662.
- Qian YH, Liang JY (2006) Rough set model based on multigranulations. In: *Proceedings of 5th IEEE Conference on Cognitive Informatics*. China I, pp 297-304.
- Qian YH, Liang JY, Yao YY, Dang CY (2010) MGRS: A multi-granulation rough set. *Inf Sci* 180:949-970.
- Qian YH, Liang JY (2011) Granulation mechanism and data modeling for complex data. Phd thesis, Shanxi University.
- Qina K, Pei Z (2005) On the topological properties of fuzzy rough sets. *Fuzzy Sets Syst* 151(3):601-613
- Qin K, Pei Z (2005) On the topological properties of fuzzy rough sets. *Fuzzy Sets Syst* 151:601-613.
- She YH, He XL (2012) On the structure of the multigranulation rough set model. *Knowl Based Syst* 36:81-92.

39. Slowinski R, Vanderpooten D (2000) A generalized definition of rough approximations based on similarity. *IEEE Trans Knowl Data Eng* 12:331-336.
40. Srivastava AK, Tiwari SP (2003) On relationships among fuzzy approximation operators, fuzzy topology, and fuzzy automata. *Fuzzy Sets Syst* 138:197-204.
41. Skowron A(1988) On the topology in information systems. *Bull Polish Academy Sciences-Mathematics* 36:477-480.
42. Xu WH, Wang QR, Zhang XT(2011) Multi-granulation fuzzy rough sets in a fuzzy tolerance approximation space. In *J Fuzzy Syst* 13(4):246-259.
43. Wang CZ, Chen DG , Hu QH (2012) On rough approximations of groups. *Int J Mach Learn Cybern*. DOI 10.1007/s13042-012-0108-6.
44. Yang XB, Song XN, Dou HL(2011) Multi-granulation rough set: from crisp to fuzzy case, *Annals of Fuzzy Math Inf* 1(1):55-70.
45. Wiweger A (1988) On topological rough sets. *Bull Polish Academy of Sciences-Mathematics* 37:51-62.
46. Wu QE, Wang T, Huang YX, Li JS (2008) Topology theory on rough sets. *IEEE Trans Syst, Man, Cybernetics-part Cybernet* 1(38):68-77.
47. Yao YY (1996) Two views of the theory of rough sets in finite universes. *Int J Approx Reason* 15:291-317.
48. Yao YY (1998) Constructive and algebraic methods of the theory of rough sets. *Inf Sci* 109:21-47.
49. Yang LY, Xu LS (2011) Topological properties of generalized approximation spaces. *Inf Sci* 182(17):3570-3580.
50. Zakowski W (1983) Approximations in the space (U, Π) . *Demonstr Mathematica* 16:761-769.
51. Zhu W (2007) Topological approaches to covering rough sets. *Inf Sci* 177(6):1499-1508.
52. Zhu W, Wang FY (2007) On three types of covering rough sets. *IEEE Trans Knowl Data Eng* 19(8):1131-1144.
53. Zhu W, Wang FY (2003) Reduction and axiomization of covering generalized rough sets. *Inf Sci* 152(1):217-230.